
Assessing the performance of a prediction error criterion model selection algorithm in the context of ARCH models

Stavros Degiannakis^{a,b,*} and Evdokia Xekalaki^{a,c}

^a*Department of Statistics, Athens University of Economics and Business, Athens 10434, Greece*

^b*Department of Informatics and Biomedicine, University of Central Greece, Lamia GR-35 100, Greece*

^c*Department of Statistics, University of California, Berkeley, CA, USA*

A number of ARCH models are considered in the framework of evaluating the performance of a method for model selection based on a standardized prediction error criterion (SPEC). According to this method, the ARCH model with the lowest sum of squared standardized forecasting errors is selected for predicting future volatility. A number of statistical criteria, that measure the distance between predicted and inter-day realized volatility, are used to examine the performance of a model to predict future volatility, for forecasting horizons ranging from one day to 100 days ahead. The results reveal that the SPEC model selection procedure has a satisfactory performance in picking that model that generates ‘better’ volatility predictions. A comparison of the SPEC algorithm with a set of other model evaluation criteria yields similar findings. It appears, therefore, that it can be regarded as a tool in guiding the choice of the appropriate model for predicting future volatility, with applications in evaluating portfolios, managing financial risk and creating speculative strategies with options.

I. Introduction

To evaluate their accuracy, volatility forecasts have to be compared with realized volatility, which cannot be observed. In the literature, it is common practice to refer the observed squared returns as the actual volatility. In this paper, a number of evaluation criteria are used to examine the ability of the SPEC model selection algorithm introduced by Degiannakis and Xekalaki (2005) to indicate the ARCH model that generates ‘better’ volatility predictions, for a

forecasting horizon ranging from one day to 100 trading days ahead.

Degiannakis and Xekalaki (2001) and Xekalaki and Degiannakis (2005) examined the performance of the SPEC algorithm through the use of economic loss functions. Degiannakis and Xekalaki (2001) made a comparative study among a set of ARCH model selection algorithms in order to examine which method yields the highest profits by trading straddles based on ten-days to 40-days-ahead variance forecasts. The results showed that the SPEC algorithm

*Corresponding author. E-mail: sdegia@aueb.gr

achieved the highest rate of return. In the context of a simulated option market, Xekalaki and Degiannakis (2005) have found that the SPEC algorithm performs better than any other comparative method of model selection in forecasting one-day-ahead conditional variance.

In this paper, evaluating the SPEC method through the implementation of statistical loss functions is considered. Specifically, the performance of the SPEC algorithm is examined through measuring the closeness of the volatility forecasts to the interday realizations. The results show that the SPEC model selection procedure has a satisfactory performance in selecting that ARCH model that tracks realized volatility closer, for a forecasting horizon ranging from 16 days to 36 days ahead. So, it is possible to use this model selection method in financial applications requiring volatility forecasts for a period longer than one day, such as option pricing or risk management. The majority of studies investigate the volatility forecasting accuracy for daily horizons, despite the fact that the practitioners require predictions of lower frequency. The Basle Committee on Banking Supervision (1998) for the use of Value-at-Risk methods requires the estimation of ten-days-ahead volatility predictions, whereas fund managers re-balance their portfolios on at least a monthly basis.

In Section II of the paper, the ARCH process is presented. Section III describes the SPEC model selection algorithm in the context of ARCH models. Section IV provides a brief description of the evaluation criteria and the interday realized volatility measures considered. In section V, the ability of the method proposed to select the ARCH model that generates 'better' predictions of the volatility, is examined. In Section VI, the proposed model selection method is compared to other methods of model selection. Finally, in Section VII, a brief discussion on the results and on the merit of looking

into the performance of the SPEC algorithm in other econometric set-ups is provided.

II. The ARCH Process

For P_t denoting the price of an asset at time t , let $y_t = \ln(P_t/P_{t-1})$ denote the continuously compounded return series of interest. The return series is decomposed into two parts, the predictable and unpredictable component:

$$y_t = E(y_{t|t-1}) + \varepsilon_t \quad (1)$$

where $E(y_{t|t-1})$ is the conditional mean of return at period t depending upon the information set available at time $t-1$ and ε_t is the prediction error. Usually, the predictable component is either the overall mean or a first order autocorrelated process (imposed by non-synchronous trading).¹ The conditional mean, unfortunately, does not have the ability to give useful predictions. That is why modern financial theory assumes the asset returns are unpredictable. Before the start of the 1980s, the view taken about returns in financial markets was that they behave as random walks and the Brock *et al.* (1987) [BDS] statistic has widely been used to test the null hypothesis that asset returns are independently and identically distributed. This hypothesis, however, has been rejected in a vast number of applications. A rejection of the null hypothesis is consistent with some types of dependence in the data, which could result in from a linear stochastic system, a non-linear stochastic system, or a non-linear deterministic system. Thus, a question arises: 'Are the non-linearities connected with the conditional mean (so, as to be used to predict future returns) or with higher order conditional moments?' Artificial neural networks,² chaotic dynamical systems,³ non-linear parametric and non-parametric models⁴ are some examples from the literature

¹ According to Campbell *et al.* (1997), 'The non-synchronous trading or non-trading effect arises when time series, usually asset prices, are taken to be recorded at time intervals of one length when in fact they are recorded at time intervals of other, possible irregular lengths.' For more details on non-synchronous trading see Scholes and Williams (1977), Dimson (1979), Cohen *et al.* (1983) and Lo and MacKinlay (1988, 1990).

² For an overview of the Neural Networks (NN) literature, see Poggio and Girosi (1990), Hertz *et al.* (1991), White (1992), Hutchinson *et al.* (1994). Plasmans *et al.* (1998) and Franses and Homelen (1998) investigated the ability of NN on forecasting exchange rates. The non-linearity found in exchange rates is due to ARCH effects. Saltoglu (2003) investigated the forecasting ability of NN on interest rates and noted the importance of modelling both the first and second moments jointly. Jasic and Wood (2004) and Perez-Rodriguez *et al.* (2005) provided evidence that NN models have a superior ability compared to other model frameworks in predicting stock indices.

³ Brock (1986), Holden (1986), Thompson and Stewart (1986) and Hsieh (1991) review applications of chaotic systems to financial markets. Adrangi and Chatrath (2003) found that the non-linearities in commodity prices are not consistent with chaos but they are explained by an ARCH process. On the other hand, Barkoulas and Travlos (1998) mentioned that even after accounting for the ARCH effect, the evidence is consistent with a chaotic structure of the Greek stock market.

⁴ Priestley (1988), Tong (1990) and Teräsvirta *et al.* (1994) cover a wide variety of non-linear models. Applications of SETAR and ARFIMA models can be found in Peel and Speight (1996) and Barkoulas *et al.* (2000), respectively.

dealing with conditional mean predictions. ARCH models⁵ and stochastic volatility models⁶ are examples from the literature dealing with conditional variance modelling. However, no non-linear models that can significantly outperform even the simplest linear model in out-of-sample forecasting seem to exist in the literature (neither in the field of stochastic non-linear models nor in the field of deterministic chaotic systems). On the other hand, the ARCH processes and stochastic volatility models appear to be more appropriate to interpret non-linearities in financial systems on the basis of the conditional variance. If an ARCH process is the true data generating mechanism, the non-linearities cannot be exploited to generate improved point predictions relative to a linear model.

In the sequel, the conditional mean is considered as an κ th order autoregressive process defined by

$$E(y_{t|t-1}) = c_0 + \sum_{i=1}^{\kappa} c_i y_{t-i} \quad (2)$$

Assuming the unpredictable component in Equation 1 is an ARCH process, it can be represented as:

$$\begin{aligned} \varepsilon_t &= z_t \sigma_t \\ z_t &\stackrel{\text{iid}}{\sim} N(0, 1) \\ \sigma_t^2 &= g(\sigma_{t-1}, \sigma_{t-2}, \dots; \varepsilon_{t-1}, \varepsilon_{t-2}, \dots; \nu_{t-1}, \nu_{t-2}, \dots), \end{aligned} \quad (3)$$

where $\{z_t\}$ is a sequence of independently and identically distributed random variables, σ_t is a time-varying, positive measurable function of the information set at time $t-1$, I_{t-1} , ν_t is a vector of predetermined variables included in I_t and $g(\cdot)$ could be a linear or non-linear functional form as is usually assumed in the ARCH literature. The unpredictable component has variance σ_t^2 , conditional on information given at time $t-1$. The conditional variance is a linear or non-linear function of lagged conditional variances, past prediction errors and predetermined variables measurable at time $t-1$. The conditional prediction error is normally distributed, but the unconditional prediction error and the conditional variance of it have an unknown form of distribution. The conditional standardized prediction error, $z_{t|t-1}$, is standard normally distributed:

$$\varepsilon_{t|t-1} \sim N(0, \sigma_t^2) \Leftrightarrow z_{t|t-1} \equiv \varepsilon_{t|t-1} \sigma_t^{-1} \sim N(0, 1) \quad (4)$$

In the recent literature, can be found a vast number of parametric specifications of ARCH models

motivated by the characteristics explored in financial markets. A researcher, who is looking for the ‘best’ model, would have in mind a variety of candidate models. The most commonly used conditional variance functions are the GARCH (Bollerslev, 1986), the Exponential GARCH, or EGARCH, (Nelson, 1991) and the Threshold GARCH, or TARARCH, (Glosten *et al.*, 1993) specifications. In the sequel, these ARCH models are considered in the following forms:

The GARCH (p,q) model

$$\sigma_t^2 = a_0 + \sum_{i=1}^q (a_i \varepsilon_{t-i}^2) + \sum_{i=1}^p (b_i \sigma_{t-i}^2) \quad (5)$$

The EGARCH (p,q) model

$$\begin{aligned} \ln(\sigma_t^2) &= a_0 + \sum_{i=1}^q \left(a_i \left| \frac{\varepsilon_{t-i}}{\sigma_{t-i}} \right| + \gamma_i \left(\frac{\varepsilon_{t-i}}{\sigma_{t-i}} \right) \right) \\ &+ \sum_{i=1}^p (b_i \ln(\sigma_{t-i}^2)) \end{aligned} \quad (6)$$

The TARARCH (p,q) model

$$\sigma_t^2 = a_0 + \sum_{i=1}^q (a_i \varepsilon_{t-i}^2) + \gamma \varepsilon_{t-1}^2 d_{t-1} + \sum_{i=1}^p (b_i \sigma_{t-i}^2) \quad (7)$$

where $d_t = 1$ if $\varepsilon_t < 0$, and $d_t = 0$ otherwise.

The majority of practical applications, i.e. option pricing, determination of the value-at-risk, require more than one-day-ahead volatility forecasts. More than one-step-ahead forecasts can be computed by repeated substitution. The forecast recursion relation of the GARCH(p,q) model is:

$$\hat{\sigma}_{t+1|t}^2 = a_0^{(t)} + \sum_{i=1}^q \left(a_i^{(t)} \varepsilon_{t-i+1}^2 \right) + \sum_{i=1}^p \left(b_i^{(t)} \sigma_{t-i+1}^2 \right) \quad (8a)$$

$$\begin{aligned} \hat{\sigma}_{t+s|t}^2 &= a_0^{(t)} + \sum_{\substack{i=1 \\ \text{for } i < s}}^q \left(a_i^{(t)} \sigma_{t-i+s}^2 \right) + \sum_{\substack{i=s \\ \text{for } i \geq s}}^q \left(a_i^{(t)} \varepsilon_{t-i+s}^2 \right) \\ &+ \sum_{i=1}^p \left(b_i^{(t)} \sigma_{t-i+s}^2 \right) \end{aligned} \quad (8b)$$

For $s > t$, the forecast of the predictive error ε_s conditional on information available at time t equals to its zero expected value, $E(\varepsilon_s | I_t) = 0$. On the other hand, the estimated value of ε_s^2 measured at time t should be equal to $\sigma_{s|t}^2$ for $s > t$. For $s \leq t$, the predictive error and its square are computed by the model with the available information at time t .

⁵ For an overview of the ARCH literature see Bollerslev *et al.* (1992, 1994), Bera and Higgins (1993), Hamilton (1994), Gouriéroux (1997) and Degiannakis and Xekalaki (2004).

⁶ For details, see Taylor (1994), Shephard (1996) and Selcuk (2005).

The forecast recursion relationship associated with the EGARCH(p, q) model is:

$$\ln(\hat{\sigma}_{t+1|t}^2) = a_0^{(i)} + \sum_{i=1}^q \left(a_i^{(i)} \left| \frac{\varepsilon_{t-i+1}}{\sigma_{t-i+1}} \right| + \gamma_i^{(i)} \left(\frac{\varepsilon_{t-i+1}}{\sigma_{t-i+1}} \right) \right) + \sum_{i=1}^p \left(b_i^{(i)} \ln(\sigma_{t-i+1}^2) \right) \quad (9a)$$

$$\ln(\hat{\sigma}_{t+s|t}^2) = a_0^{(i)} + \sum_{\substack{i=s \\ \text{for } i \geq s}}^q \left(a_i^{(i)} \left| \frac{\varepsilon_{t-i+s}}{\sigma_{t-i+s}} \right| + \gamma_i^{(i)} \left(\frac{\varepsilon_{t-i+s}}{\sigma_{t-i+s}} \right) \right) + \left(\frac{2}{\pi} \right)^{1/2} \sum_{\substack{i=1 \\ \text{for } i < s}}^q (a_i^{(i)}) + \sum_{i=1}^p \left(b_i^{(i)} \ln(\sigma_{t-i+s}^2) \right) \quad (9b)$$

that associated with the TARCh(p, q) model is:

$$\hat{\sigma}_{t+1|t}^2 = a_0^{(i)} + \sum_{i=1}^q \left(a_i^{(i)} \varepsilon_{t-i+1}^2 \right) + \gamma^{(i)} \varepsilon_t^2 d_t + \sum_{i=1}^p \left(b_i^{(i)} \sigma_{t-i+1}^2 \right) \quad (10a)$$

$$\hat{\sigma}_{t+s|t}^2 = a_0^{(i)} + \sum_{\substack{i=1 \\ \text{for } i < s}}^q \left(a_i^{(i)} \sigma_{t-i+s}^2 \right) + \sum_{\substack{i=1 \\ \text{for } i \geq s}}^q \left(a_i^{(i)} \varepsilon_{t-i+s}^2 \right) + \gamma^{(i)} \sigma_{t-1+s}^2 E(d_t) + \sum_{i=1}^p \left(b_i^{(i)} \sigma_{t-i+s}^2 \right) \quad (10b)$$

Here, $E(d_t)$ denotes the percentage of negative innovations out of all innovations. Under the assumption of normally distributed innovations, the

pairwise comparisons of the sums of squared standardized one-step-ahead forecasting errors of a set of ARCH models is provided. Degiannakis and Xekalaki (2005) introduced the SPEC model selection method based on the correlated gamma ratio (CGR) distribution, which was derived by Xekalaki *et al.* (2003) as the distribution of the ratio of two variables jointly distributed according to Kibble's (1941) bivariate gamma distribution. Kibble (1941) proves that if, for $t=1, 2, \dots$, the joint distribution of $(r_t^{(A)}, r_t^{(B)})$ is the bivariate standard normal, then the joint distribution of $2^{-1} \sum_{i=1}^T r_i^{2(A)}$ and $2^{-1} \sum_{i=1}^T r_i^{2(B)}$ is Kibble's bivariate gamma distribution. As pointed out by Xekalaki *et al.* (2003), $r_t^{(A)}$ and $r_t^{(B)}$ could represent the standardized prediction errors from two regression models (not necessarily nested) but with a common dependent variable. The distribution of the ratio of the sum of their squares is the CGR distribution; symbolically, $\sum_{i=1}^T r_i^{2(B)} / \sum_{i=1}^T r_i^{2(A)} \sim CGR(k, \rho)$, where $k = T/2$ and $\rho = Cor(r_t^{(A)}, r_t^{(B)})$. Thus, two regression models can be compared through testing the null hypothesis of equivalence of the models in their predictability against the alternative that model (A) produces 'better' predictions. The null hypothesis is rejected if $\sum_{i=1}^T r_i^{2(B)} / \sum_{i=1}^T r_i^{2(A)} > CGR(k, \rho, \alpha)$, where $CGR(k, \rho, \alpha)$ is the $100(1 - \alpha)$ percentile of the CGR distribution.⁷

Degiannakis and Xekalaki (2005) assumed that interest is in comparing the predictive ability of two ARCH models:

Model A	Model B
$\varepsilon_i^{(A)} = z_{1,t} \sigma_i^{(A)}$	$\varepsilon_i^{(B)} = z_{2,t} \sigma_i^{(B)}$
$z_{1,t} \stackrel{iid}{\sim} N(0, 1)$	$z_{2,t} \stackrel{iid}{\sim} N(0, 1)$
$\sigma_i^{2(A)} = g\left(\sigma_{i-1}^{2(A)}, \dots, \sigma_{i-p}^{2(A)}, \varepsilon_{i-1}^{2(A)}, \dots, \varepsilon_{i-q}^{2(A)}, \nu_{i-1}^{(A)}, \nu_{i-2}^{(A)}, \dots\right)$	$\sigma_i^{2(B)} = g\left(\sigma_{i-1}^{2(B)}, \dots, \sigma_{i-p}^{2(B)}, \varepsilon_{i-1}^{2(B)}, \dots, \varepsilon_{i-q}^{2(B)}, \nu_{i-1}^{(B)}, \nu_{i-2}^{(B)}, \dots\right)$

expected number of negative shocks is equal to the expected number of positive shocks, or $E(d_t) = 0.5$.

The forecast of the conditional variance at time t over a horizon of N days ahead is simply the average of the estimated future variance conditional on information given at time t

$$\sigma_{t(N)}^2 = N^{-1} \sum_{i=1}^N \hat{\sigma}_{t+i|t}^2 \quad (11)$$

III. The SPEC Model Selection Method

In this section, a brief description of the theoretical motivation of the SPEC algorithm that is based on

The joint distribution of $2^{-1} \sum_{i=1}^T \hat{z}_{i|t-1}^{2(A)} \equiv 2^{-1} \sum_{i=1}^T \hat{\varepsilon}_{i|t-1}^{2(A)} / \hat{\sigma}_{i|t-1}^2$ and $2^{-1} \sum_{i=1}^T \hat{z}_{i|t-1}^{2(B)} \equiv 2^{-1} \sum_{i=1}^T \hat{\varepsilon}_{i|t-1}^{2(B)} / \hat{\sigma}_{i|t-1}^2$ is Kibble's bivariate gamma distribution. Thus, the standardized one-step-ahead prediction errors can be used to test the null hypothesis of equivalence of the models in their predictive ability against the alternative that the first model produces 'better' predictions. The null hypothesis is rejected if $\sum_{i=1}^T \hat{z}_{i|t-1}^{2(B)} / \sum_{i=1}^T \hat{z}_{i|t-1}^{2(A)} > CGR(k, \rho, \alpha)$. Note that the SPEC algorithm is always computed on the basis of the one-step-ahead forecasts since $\hat{z}_{t+1|t}$ are asymptotically normally distributed (Degiannakis and Xekalaki, 2005), while the standardized

⁷Percentage points of the CGR distribution can be found in Xekalaki *et al.* (2003) and Degiannakis and Xekalaki (2005).

Table 1. The estimation steps required at time k for each model m by the SPEC model selection algorithm. At time k ($k = T, T + 1, \dots$), select the model m with the minimum value for the sum of the squares of the T most recent standardized one-step-ahead prediction errors

$$\sum_{t=k-T+1}^k \hat{z}_t^{2(m)} \equiv \sum_{t=k-T+1}^k \hat{\varepsilon}_{t|t-1}^{2(m)} / \hat{\sigma}_{t|t-1}^{2(m)}$$

Time					
Model	$k = T$	$k = T + 1$...	$k = T + K$...
$m = 1$	$\sum_{t=1}^T \hat{z}_{t t-1}^{2(1)}$	$\sum_{t=2}^T \hat{z}_{t t-1}^{2(1)}$...	$\sum_{t=K+1}^{T+K} \hat{z}_{t t-1}^{2(1)}$...
$m = 2$	$\sum_{t=1}^T \hat{z}_{t t-1}^{2(2)}$	$\sum_{t=2}^{T+1} \hat{z}_{t t-1}^{2(2)}$...	$\sum_{t=K+1}^{T+K} \hat{z}_{t t-1}^{2(2)}$...
\vdots	\vdots	\vdots	\ddots	\vdots	\vdots
$m = M$	$\sum_{t=1}^T \hat{z}_{t t-1}^{2(M)}$	$\sum_{t=2}^{T+1} \hat{z}_{t t-1}^{2(M)}$...	$\sum_{t=K+1}^{T+K} \hat{z}_{t t-1}^{2(M)}$...

residuals from N -step ahead forecasts, $\hat{z}_{t+N|t}$, for $N \geq 2$, are not.

According to the SPEC model selection algorithm, the models that are considered as having a ‘better’ ability to predict future volatility of the dependent variable, are those with the lowest sum of squared standardized one-step-ahead prediction errors. Assume that M candidate ARCH models are available and that the ‘most suitable’ model at each of a sequence of points in time is being looked for. On the basis of the SPEC algorithm, at time k , selecting a strategy for the most appropriate model to forecast volatility at time $k + 1$ ($k = T, T + 1, \dots$) could naturally amount to selecting the model which, at time k , has the lowest sum of squared standardized one-step-ahead prediction errors. So, each time the SPEC model selection method is applied, the model used to predict the conditional variance is revised. Table 1 summarizes the estimation steps comprising this approach. On the face of it, the view might be taken that a model can always be made more attractive simply through over-predicting the volatility. However, an algorithm constructed so as to select the model with the maximum sum of the T most recent estimated one-step-ahead volatility forecasts will not pick the same models as those picked by the SPEC model selection method.

In the next section, the methodology applied to evaluate the performance of a model in estimating

future volatility is presented, while in Section V, the ability of the SPEC model selection algorithm to indicate those ARCH models that generate ‘better’ volatility predictions is illustrated on daily returns of the S&P500 stock index.

IV. Evaluating the Volatility Forecast Performance

The main problem in evaluating the predictive performance of a model is the choice of the function to be used to measure the distance between estimations and observations. Evaluating the performance of the variance forecasts requires knowledge of the actual volatility, which is unobservable. Thus, in evaluating the predictive performance of a variance model a question of a dual nature arises: that of determining the realized volatility and of considering the appropriate measure to evaluate the closeness of the forecasts to the corresponding realizations.

Realized volatility measures

Practitioners’ most popular volatility measures are the average of squared daily returns and the variance of the daily returns. These measures, expressed on a daily basis for a horizon of N days ahead, are:

$$s_{i(N)}^2 = N^{-1} \sum_{i=1}^N y_{t+i}^2 \tag{12}$$

$$\bar{s}_{i(N)}^2 = (N - 1)^{-1} \sum_{i=1}^N (y_{t+i} - \bar{y}_{i(N)})^2 \tag{13}$$

respectively, where $\bar{y}_{i(N)} = N^{-1} \sum_{i=1}^N y_{t+i}$ is the average return. The interday volatility measures are the most popular measures. However, as noted in the literature (e.g. Ebens, 1999), although the squared daily returns are unbiased volatility estimators, they are very noisy. Note that, under the ARCH process, the squared return can be represented by $y_t^2 = z_t^2 \sigma_t^2$. It is therefore defined as the product of the true volatility times the square of a normally distributed process. Recently, Alizadeh *et al.* (2002) and Sadorsky (2005) proposed the log range measure of volatility defined as the difference between the highest and lowest log asset prices over the interval of N days. In the present paper, the popular among practitioners inter-day measures are utilized. An investigation

based on the intraday realized volatility is worth future exploration.⁸

Evaluation criteria

A large number of forecast evaluation criteria exists in the literature. However, none is generally acceptable. Because of high non-linearity in volatility models and the variety of statistical evaluation criteria, a number of researchers constructed economic criteria based upon the goals of their particular application. West *et al.* (1993) develop a criterion based on the decisions of a risk averse investor. Engle *et al.* (1993) assume that the objective is to price options and develop a loss function from the profitability of a particular trading strategy. González-Rivera *et al.* (2004) considered comparing the performance of various volatility models on the basis of economic and statistical loss functions. Their study revealed that there does not exist a unique model that can be regarded as the best performer across various loss functions. Brooks and Persaud (2003) also found that the forecasting accuracy of various methods considered in the literature is highly sensitive to the measure used to evaluate them. Hence, different loss functions may point towards different models as the most appropriate in volatility forecasting. In the sequel, focus is on statistical criteria to measure the closeness of the forecasts to the realizations, in order to avoid restrictions imposed by economic theory. Moreover, statistical criteria are considered that are robust to non-linearity and heteroscedasticity. Pagan and Schwert (1990) use statistical criteria to compare the in-sample and out-of-sample performance of parametric and non-parametric ARCH models. Besides, Heynen and Kat (1994) investigate the predictive performance of ARCH and stochastic volatility models and Hol and Koopman (2000) compare the predictive ability of stochastic volatility and implied volatility models. Andersen *et al.* (1999) applied heteroscedasticity-adjusted statistics to examine the forecasting performance of intraday returns. Denoting the forecasting variance over an N day period measured at day t by $\sigma_{t(N)}^2$, and the realized variance over the same period by $s_{t(N)}^2$, the following evaluation criteria are considered:

$$\text{Squared Error (SE): } \left(\sigma_{t(N)}^2 - s_{t(N)}^2 \right)^2 \quad (14)$$

$$\text{Absolute Error (AE): } \left| \sigma_{t(N)}^2 - s_{t(N)}^2 \right| \quad (15)$$

Heteroscedasticity Adjusted Squared Error (HASE):

$$\left(1 - \frac{s_{t(N)}^2}{\sigma_{t(N)}^2} \right)^2 \quad (16)$$

Heteroscedasticity Adjusted Absolute Error (HAAE):

$$\left| 1 - \frac{s_{t(N)}^2}{\sigma_{t(N)}^2} \right| \quad (17)$$

Logarithmic Error (LE):

$$\ln \left(\frac{s_{t(N)}^2}{\sigma_{t(N)}^2} \right)^2 \quad (18)$$

The first two functions have been widely used in the literature (see, e.g. Heynen and Kat, 1994; West and Cho, 1995; Yu, 2002; Brooks and Persaud, 2003). The HASE and HAAE functions were considered by Walsh and Tsou (1998) and Andersen *et al.* (1999), while the LE function was utilized by Pagan and Schwert (1990) and Saez (1997).

Usually, the average of the evaluation criteria is computed. However, when simulating an AR(1) GARCH(1,1) process, which is the most commonly used model in financial applications, the distributions of $(\sigma_{t(N)}^2 - s_{t(N)}^2)$, $(1 - s_{t(N)}^2/\sigma_{t(N)}^2)$ and $\ln(s_{t(N)}^2/\sigma_{t(N)}^2)$ are asymmetric with extreme outliers. It would therefore be advisable to compute both the mean and the median of the evaluation criteria. Figure 1 depicts the histograms of the one-step forecast error distribution from the following simulated process:

$$\begin{aligned} y_t &= 0.001 + 0.1y_{t-1} + \varepsilon_t \\ \sigma_t^2 &= 0.002 + 0.05\varepsilon_{t-1}^2 + 0.9\sigma_{t-1}^2 \\ \varepsilon_t &= \sigma_t z_t \quad z_t \stackrel{\text{iid}}{\sim} N(0, 1). \end{aligned} \quad (19)$$

V. Examining the Performance of the SPEC Model Selection Algorithm

In this section, the ability of the SPEC model selection algorithm to lead to the ARCH models that track closer future volatility is illustrated on a series of daily log-returns. As follows from Section II, the return series can be modelled in the following form:

$$\begin{aligned} y_t &= c_0 + \sum_{i=1}^k c_i y_{t-i} + \varepsilon_t \\ \varepsilon_t &= z_t \sigma_t \quad z_t \stackrel{\text{iid}}{\sim} N(0, 1) \\ \sigma_t^2 &= g(\sigma_{t-1}, \sigma_{t-2}, \dots; \varepsilon_{t-1}, \varepsilon_{t-2}, \dots; \nu_{t-1}, \nu_{t-2}, \dots) \end{aligned} \quad (20)$$

⁸ For details and references about intra-day realized volatility the interested reader is referred to Andersen and Bollerslev (1997, 1998a, b), Barndorff-Nielsen and Shephard (1998), Andersen *et al.* (1999, 2000a, b, 2001a, b, 2003, 2005).




Fig. 1(a). Histogram of $\hat{\sigma}_{t+1|t}^2 - y_{t+1}^2$ from an AR(1)GARCH(1,1) simulated process, (b) Histogram of $1 - y_{t+1}^2 / \hat{\sigma}_{t+1|t}^2$ from an AR(1)GARCH(1,1) simulated process, (c) Histogram of $\ln(y_{t+1}^2 / \hat{\sigma}_{t+1|t}^2)$ from an AR(1)GARCH(1,1) simulated process

In the sequel, the above form is considered in connection with the ARCH models defined by Equations 5–7, for $\kappa = 0, 1, 2, 3, 4$, $p = 0, 1, 2$ and $q = 1, 2$, thus yielding a total of 85 cases.⁹ The first autoregressive order of the conditional mean accounts for the non-synchronous trading. However, various orders are considered as, according to Hansen and Lunde (2003), a small improvement of the modelling of the conditional mean, may lead to a clear improvement in the forecast of volatility. Also, contrary to the majority of applied studies, such as those by Klaassen (2002), Vilasuso (2002) and Hansen and Lunde (2003), where the forecasts are calculated by estimating the models once, in the present study the models are re-estimated every trading day, in order to evaluate the forecast accuracy under real world circumstances. Further, as Degiannakis and Xekalaki’s (2005) results reveal, the SPEC algorithm can be applied to all the conditional variance functions with consistent and asymptotically normal estimators of the parameters. Thus, although, austere mathematically, minimizing the sum of squared standardized one-step-ahead residuals is not equivalent to maximizing the likelihood, Bollerslev and Wooldridge’s (1992)

quasi-maximum likelihood method is employed in the sequel to estimate the models. Maximum likelihood estimates of the parameters are obtained by numerical maximization of the log-likelihood function using the Marquardt algorithm (Marquardt, 1963), a modification of the Berndt, Hall, Hall and Hausman, or BHHH, algorithm (Berndt *et al.*, 1974). The quasi-maximum likelihood estimator (QMLE) is used, as according to Bollerslev and Wooldridge, it is generally consistent, has a normal limiting distribution and provides asymptotic standard errors that are valid under non-normality.

The data set consists of 1661 S&P500 daily log-returns in the period from 24 November 1993 to 26 June 2000. Although, large data sets are often used in the literature for the estimation of ARCH models, here using a not too large sample is considered, which would expectantly incorporate changes in trading behaviour more efficiently as the evidence is from various findings in the literature (e.g. Engle *et al.*, 1993; Frey and Michaud, 1997; Angelidis *et al.*, 2004; Xekalaki and Degiannakis, 2005). So, the ARCH processes are estimated using a rolling sample of constant size equal to 500. Thus, the first one-step-ahead volatility prediction, $\hat{\sigma}_{t+1|t}^2$, is available at

⁹ Numerical maximization of the log-likelihood function, for the EGARCH(2,2) model, failed to converge in more than 1% of the trading days. So the five EGARCH models for $p = q = 2$ were excluded.

time $t = 500$. Applying the SPEC model selection algorithm, the sum of squared standardized one-step-ahead prediction errors, $\sum_{t=1}^T \hat{z}_{t|t-1}^2$, was estimated considering various values for T , and, in particular, $T = 5(5)80$.¹⁰ Thus, the evaluation criteria were applied on the one-step-ahead forecasts using $1661 - 500 - 80 = 1081$ data points, on the two-step-ahead forecasts using $1661 - 500 - 81 = 1080$ data points, . . . , and on the N th-step-ahead forecasts using $1081 - N + 1$ data points.

Adopting Brooks and Persaud's (2003) approach evaluating multistep-ahead forecasts based on overlapping time periods are considered. In particular, most of the studies in the literature evaluate the multistep forecasts using non-overlapping time periods in order to infer about the statistical significance of the ranking. The main purpose is to examine the application potential of the SPEC algorithm of selection of models on the basis of their forecasting ability in terms of volatility. So, the mean and the median value of each of the five evaluation criteria, in Equations 14–18, were computed, yielding a total of ten evaluation criteria for each forecasting horizon from one day to 100 days ahead. However, volatility is expressed either as the variance or as the standard deviation. Thus, in order to examine possible differences between forecasting the variance and its square root, the evaluation criteria were, also, applied on the standard deviation. Therefore, $\sigma_{t(N)}^2$ and $s_{t(N)}^2$, in Equations 14–18, were replaced by $\sigma_{t(N)}$ and $s_{t(N)}$, respectively and ten more evaluation criteria were computed. In total, 20 evaluation criteria were computed for a horizon ranging from one trading day to five trading months. In Section IV, two realized volatility measures were mentioned. As, qualitatively, they are of the same nature, in the sequel, the analysis is based on the realized volatility as defined by $s_{t(N)}^2$.¹¹

It was examined whether the ARCH models selected by the SPEC algorithm achieve the lowest value of the evaluation criteria. The main focus was on the median values of the criteria and mainly on the heteroscedasticity adjusted criteria since they are more robust to asymmetry. The comparative evaluation is performed by computing the loss functions for variance forecasts always obtained by a single model on the one hand, and for variance forecasts obtained by models picked by the SPEC algorithm on the other. Table 2, presents the ARCH model that achieved the minimum value of each evaluation criterion. Table 2 refers to a subset of the forecasting horizon, but it is representative for the total set of

100 trading days ahead. The SPEC algorithm is applied for 16 values for T , and, in particular, $T = 5(5)80$. The SPEC(T) value refers to the size T for which the SPEC algorithm achieves the minimum value of the evaluation criteria. The minARCH and minSPEC values refer to the minima of the evaluation criteria achieved by a single model and by models picked by the SPEC algorithm, respectively. As concerns the variance forecasts obtained by any single model, the results are in line with those existing in the literature, i.e. they are not consistent across all functions. Although, the exponential ARCH specification exhibits the best performance in the majority of the cases (84.1% of the cases presented in Table 2), the autoregressive order of the conditional mean is not constant across the evaluation criteria. However, the lag order $p = q = 1$ of the EGARCH variance specification exhibits the best performance in 63.4% of the cases.

Figure 2 shows, for each evaluation criterion and each forecasting horizon, whether ARCH models selected by the SPEC algorithm achieve the lowest value of the evaluation criteria. In the first part of Fig. 2, the performance of the models, which are selected by the SPEC algorithm, on the basis of the conditional variance is depicted, while, the second part refers to their performance on forecasting standard deviation. The general conclusion is that the SPEC algorithm leads to the selection of the ARCH processes which track closer the realized volatility in the majority of the cases. Specifically, for the forecasting horizon ranging from 11 to 52 days, the models selected by the SPEC algorithm achieve the lowest criteria values, irrespectively of the evaluation criteria. The percentage of cases, in which the models picked by the SPEC algorithm achieve the lowest value of the evaluation criteria, is higher around the forecasting horizon ranging from 16 to 36 days ahead, or 4–7 trading weeks ahead. The result is in accordance to Degiannakis and Xekalaki (2001) who provided evidence that option's traders using variance forecasts for horizons ranging from 10 to 40 trading days obtained by models suggested by the SPEC algorithm achieved the highest rate of return among a set of model selection criteria. Table 3 presents the percentage of cases the models selected by the SPEC algorithm perform 'better' than any other single model as judged by the evaluation criteria, for three different horizon ranges. Note that, in terms of the MSE and MAE criteria, none of the models chosen by the SPEC algorithm appears to perform better in any of the forecasting

¹⁰ Here, $T = a(b)c$ denotes $T = a, a + b, a + 2b, \dots, c - b, c$.

¹¹ The analysis was also conducted based on $\hat{s}_{t(N)}$ giving qualitatively similar results.

Table 2. Minimum values of the evaluation criteria for variance forecasts always obtained by a single ARCH model on the one hand and for variance forecasts obtained by models suggested by the SPEC algorithm on the other, for a subset of the forecasting horizon which ranges from 5 to 100 days. The first panel refers to the variance, ($k=2$), whereas the second panel accounts for the standard deviation, ($k=1$). Model refers to the single ARCH model that achieves the lowest value of the evaluation criterion. The SPEC(T) value refers to the size T for which the SPEC algorithm achieves the minimum value of the evaluation criteria. The minARCH and minSPEC values refer to the minima of the evaluation criteria achieved by a single ARCH model and by models suggested by the SPEC algorithm, respectively. The evaluation criteria are the annualized mean and median values of the following loss functions:

Squared error (SE): $(\sigma_{i(t)}^k - s_{i(t)}^k)^2$
 Absolute error (AE): $|\sigma_{i(t)}^k - s_{i(t)}^k|$
 Heteroscedasticity adjusted squared error (HASE): $(1 - s_{i(t)}^k / \sigma_{i(t)}^k)^2$
 Heteroscedasticity adjusted absolute error (HAAE): $|1 - s_{i(t)}^k / \sigma_{i(t)}^k|$
 Logarithmic error (LE): $\ln(s_{i(t)}^k / \sigma_{i(t)}^k)^2$

		Forecast horizon in days																							
		5	10	15	16	18	20	22	24	26	28	30	32	34	35	40	45	50	60	70	80	90	100		
Evaluation criteria for the conditional variance, $k=2$																									
MeanSE																									
Model	AR(4)	AR(4)	AR(4)	AR(4)	AR(4)	AR(4)	AR(4)	AR(4)	AR(4)	AR(3)	AR(4)	AR(4)	AR(4)	AR(4)	AR(4)	AR(4)	AR(4)	AR(4)	AR(4)	AR(2)	AR(1)	AR(1)	AR(1)	AR(1)	AR(1)
minARCH	EG(1,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)
SPEC(7)	55	55	25	20	20	20	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15
minSPEC	0.187	0.126	0.097	0.092	0.083	0.077	0.072	0.069	0.067	0.065	0.064	0.063	0.062	0.061	0.059	0.056	0.053	0.049	0.045	0.041	0.038	0.031	0.030	0.030	
MeanAE																									
Model	AR(2)	AR(2)	AR(4)	AR(2)	AR(2)	AR(2)	AR(2)	AR(2)	AR(4)	AR(3)	AR(3)	AR(4)	AR(4)	AR(4)	AR(4)	AR(4)	AR(4)	AR(4)	AR(4)	AR(2)	AR(2)	AR(2)	AR(1)	AR(1)	AR(1)
minARCH	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(2,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)
SPEC(7)	25	55	25	25	20	20	20	15	15	15	15	15	15	15	15	15	15	15	15	20	20	20	20	20	20
minSPEC	1.964	1.683	1.541	1.513	1.458	1.423	1.402	1.395	1.386	1.378	1.366	1.360	1.357	1.355	1.356	1.353	1.351	1.340	1.329	1.317	1.305	1.291	1.283	1.201	
MedSE																									
Model	AR(4)	AR(4)	AR(4)	AR(4)	AR(4)	AR(4)	AR(2)	AR(2)	AR(2)	AR(2)	AR(2)	AR(2)	AR(2)	AR(2)	AR(2)	AR(2)	AR(2)	AR(2)	AR(2)	AR(2)	AR(2)	AR(0)	AR(0)	AR(0)	AR(4)
minARCH	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(1,1)	EG(0,2)	EG(0,2)	EG(0,2)	EG(0,2)
SPEC(7)	35	30	25	35	35	35	35	35	35	35	35	35	35	35	35	35	35	35	35	35	35	25	25	25	20
minSPEC	0.0119	0.0077	0.0066	0.0064	0.0055	0.0052	0.0051	0.0050	0.0050	0.0048	0.0047	0.0047	0.0045	0.0044	0.0043	0.0042	0.0041	0.0041	0.0041	0.0041	0.0040	0.0035	0.0030	0.0025	0.0056
MedAE																									
minARCH	0.970	0.824	0.784	0.751	0.770	0.755	0.740	0.748	0.722	0.703	0.696	0.690	0.674	0.682	0.682	0.666	0.653	0.657	0.650	0.650	0.694	0.690	0.671	0.671	
SPEC(7)	35	30	25	35	35	35	35	35	35	35	35	35	35	35	35	35	35	35	35	35	35	25	25	20	
minSPEC	1.089	0.877	0.815	0.799	0.742	0.723	0.717	0.708	0.708	0.690	0.684	0.683	0.668	0.658	0.658	0.683	0.661	0.642	0.672	0.706	0.729	0.749	0.749		
MeanHASE																									
Model	AR(1)	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(4)	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(1)	AR(1)	AR(1)	AR(1)	AR(1)	AR(1)	AR(1)	AR(3)
minARCH	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,2)	G(1,2)	T(1,2)	T(1,2)	T(1,2)	T(1,2)	T(1,2)	T(1,2)	G(1,2)	
SPEC(7)	55	40	20	20	20	20	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	10	
minSPEC	208.9	146.0	111.0	104.7	93.2	85.1	78.9	75.2	73.3	71.3	70.3	69.7	68.5	68.2	65.9	63.4	61.7	62.4	61.3	59.1	57.3	57.5	57.5		
MeanHAAE																									
Model	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(3)	AR(4)	AR(4)	AR(1)	AR(1)	AR(1)	AR(1)	AR(1)	AR(1)	AR(1)	AR(1)
minARCH	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	G(1,1)	T(1,2)	T(1,2)	T(1,2)	T(1,2)	T(1,2)	T(1,2)	EG(1,1)	
SPEC(7)	20	35	35	20	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	10	
minSPEC	64.52	56.37	52.23	51.22	49.53	48.15	47.53	47.37	47.38	47.33	47.15	47.32	47.51	47.68	47.99	48.49	49.09	50.71	51.74	52.64	53.31	54.09	54.09		

(continued)

Table 2. Continued

		Forecast horizon in days																						
		5	10	15	16	18	20	22	24	26	28	30	32	34	35	40	45	50	60	70	80	90	100	
MedHASE																								
Model	AR(1) EG(2,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	
minARCH	19.53	12.50	10.08	9.82	9.69	9.02	9.24	9.43	9.43	9.22	8.51	8.85	8.42	8.43	8.86	8.45	8.40	8.72	8.36	8.57	8.38	8.80	8.33	8.33
SPEC(T)	35	35	35	25	25	25	25	15	15	15	15	15	15	15	15	40	40	35	35	35	25	25	25	20
minSPEC	19.52	12.27	10.81	10.01	9.10	8.42	8.45	8.14	8.14	7.95	7.56	7.77	7.54	7.44	7.43	7.80	7.38	8.08	8.41	8.22	9.39	8.33	8.33	8.74
MedHAAE																								
minARCH	44.20	35.35	31.75	31.33	31.12	30.04	30.39	30.70	30.70	30.37	29.17	29.74	29.02	29.03	29.77	29.07	28.97	29.52	28.91	29.28	28.95	29.67	28.86	28.86
SPEC(T)	35	35	35	25	25	25	15	15	15	15	15	15	15	15	15	40	40	35	35	35	25	25	25	20
minSPEC	44.19	35.03	32.88	31.64	30.17	29.02	29.07	28.54	28.54	28.19	27.49	27.88	27.46	27.27	27.25	27.93	27.17	28.43	29.01	28.67	30.65	28.87	28.87	29.57
MeanLE																								
Model	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)
minARCH	63.02	40.18	33.82	33.10	31.36	29.90	29.03	28.31	28.31	27.70	27.19	26.84	26.42	26.21	26.06	25.40	24.77	24.41	23.99	23.40	23.01	22.44	22.44	22.27
SPEC(T)	35	40	30	30	20	20	20	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15
minSPEC	68.24	39.93	31.75	30.99	29.04	27.71	27.07	26.43	26.43	26.07	25.85	25.65	25.40	25.24	25.16	24.94	24.85	24.84	25.09	25.16	25.22	25.22	25.22	25.55
MedLE																								
Model	AR(3) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)	AR(4) EG(1,1)
minARCH	21.60	13.53	11.27	11.10	10.41	9.46	9.86	10.03	10.03	9.81	9.32	9.14	8.65	8.92	9.11	9.07	8.68	9.19	8.61	9.34	9.46	9.22	9.22	8.83
SPEC(T)	25	35	25	25	25	25	15	15	15	15	15	15	15	15	15	40	35	35	35	35	35	25	25	25
minSPEC	22.80	14.22	10.94	10.65	10.04	8.65	9.27	8.49	8.49	8.02	8.19	7.59	7.72	8.17	8.33	8.36	8.20	9.10	9.06	9.62	9.73	9.52	9.52	9.14
Evaluation criteria for the conditional standard deviation, $k=1$																								
MeanSE																								
Model	AR(2) EG(1,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)	AR(4) EG(2,1)
minARCH	0.507	0.396	0.343	0.333	0.316	0.301	0.292	0.283	0.283	0.277	0.271	0.268	0.265	0.262	0.261	0.252	0.242	0.235	0.224	0.214	0.206	0.196	0.196	0.191
SPEC(T)	55	55	30	20	20	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15
minSPEC	0.535	0.398	0.335	0.323	0.301	0.288	0.277	0.271	0.271	0.267	0.265	0.263	0.260	0.259	0.259	0.255	0.250	0.245	0.239	0.234	0.227	0.220	0.220	0.216
MeanAE																								
Model	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)
minARCH	4.697	4.001	3.766	3.718	3.632	3.576	3.543	3.535	3.535	3.515	3.487	3.465	3.432	3.430	3.423	3.401	3.360	3.356	3.338	3.315	3.327	3.308	3.308	3.284
SPEC(T)	55	55	30	25	20	20	15	15	15	15	15	15	15	15	15	15	15	15	15	15	20	20	20	20
minSPEC	4.944	4.111	3.759	3.698	3.583	3.504	3.466	3.448	3.448	3.431	3.416	3.388	3.381	3.386	3.393	3.394	3.410	3.433	3.464	3.466	3.484	3.493	3.493	3.491
MedSE																								
Model	AR(4) EG(1,1)	AR(0) EG(1,1)	AR(1) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)	AR(2) EG(1,1)
minARCH	0.1211	0.0783	0.0653	0.0632	0.0624	0.0596	0.0573	0.0575	0.0575	0.0565	0.0536	0.0548	0.0499	0.0507	0.0523	0.0500	0.0516	0.0518	0.0500	0.0506	0.0531	0.0516	0.0516	0.0486
SPEC(T)	60	35	35	35	35	35	35	35	35	15	15	20	15	20	15	20	35	35	35	35	35	35	35	35
minSPEC	0.1322	0.0823	0.0739	0.0699	0.0597	0.0551	0.0560	0.0549	0.0549	0.0538	0.0521	0.0499	0.0480	0.0474	0.0478	0.0498	0.0470	0.0490	0.0515	0.0531	0.0538	0.0542	0.0542	0.0536
MedAE																								
minARCH	3.479	2.798	2.555	2.514	2.499	2.441	2.394	2.398	2.398	2.377	2.316	2.340	2.234	2.252	2.286	2.236	2.272	2.275	2.235	2.250	2.305	2.271	2.271	2.204
SPEC(T)	60	35	35	35	35	35	35	35	35	15	15	20	15	20	15	20	35	35	35	35	35	35	35	35
minSPEC	3.636	2.868	2.719	2.645	2.444	2.347	2.367	2.344	2.344	2.319	2.282	2.234	2.190	2.177	2.186	2.232	2.168	2.213	2.269	2.304	2.319	2.327	2.327	2.315




Fig. 2(a). The plots indicate whether the ARCH models selected by the SPEC algorithm achieve the lowest value of the evaluation criterion, for a forecasting horizon ranging from one day to 100 days ahead. The value 2 indicates that the ARCH model selected by the SPEC algorithm achieves the lowest value for the corresponding criterion and the specific forecasting horizon. The value 1 indicates the opposite. The realized volatility measure is expressed as in Equation 12. The evaluation criteria are the mean and the median values of the functions defined by Equations 14–18




Fig. 2(b). The plots indicate whether the ARCH models selected by the SPEC algorithm achieve the lowest value of the evaluation criterion, for a forecasting horizon ranging from one day to 100 days ahead. The value 2 indicates that the ARCH model selected by the SPEC algorithm achieves the lowest value for the corresponding criterion and the specific forecasting horizon. The value 1 indicates the opposite. The realized volatility measure is expressed as the square root of Equation 12. The evaluation criteria are the mean and the median values of the functions defined by Equations 14–18

Table 3. The percentage of times the ARCH models selected by the SPEC algorithm perform 'better' than any other single model as judged by the evaluation criteria. The first and the second panel correspond to the mean and the median of the evaluation criteria, respectively. The left and the right part of the panels correspond to the volatility expressed as the variance and the standard deviation of the returns, respectively.

		Mean									
Days ahead forecasting horizon	Variance	Variance			Standard deviation						
		MSE	MAE	MHASE	MHAAE	MLE	MSE	MAE	MHASE	MHAAE	MLE
1-100		0%	0%	47%	56%	34%	26%	26%	54%	56%	34%
11-52		0%	0%	88%	100%	79%	62%	62%	100%	100%	79%
16-36		0%	0%	100%	100%	100%	100%	100%	100%	100%	100%
	Median										
Days ahead forecasting horizon	Variance	Variance			Standard deviation						
		MedSE	MedAE	MedHASE	MedHAAE	MedLE	MedSE	MedAE	MedHASE	MedHAAE	MedLE
1-100		40%	40%	65%	65%	35%	38%	38%	50%	50%	35%
11-52		64%	64%	88%	88%	83%	81%	81%	93%	93%	83%
16-36		86%	86%	95%	95%	100%	90%	90%	100%	100%	100%

Notes: MSE: Mean square error.

MAE: Mean absolute error.

MHASE: Mean heteroscedasticity adjusted squared error.

MHAAE: Mean heteroscedasticity adjusted absolute error.

MLE: Mean logarithmic error.

MedSE: Median square error.

MedAE: Median absolute error

MedHASE: Median heteroscedasticity adjusted squared error.

MedHAAE: Median heteroscedasticity adjusted absolute error.

MedLE: Median logarithmic error.




Fig. 3. Sample size of the SPEC model selection algorithm, suggested by the evaluation criteria

horizons considered. But, in terms of the median values of the criteria and the heteroscedasticity adjusted criteria, which are robust to asymmetry, the models selected by the SPEC algorithm appear to have a better performance than any other single model in all the forecasting horizons.

It is interesting to note that, via the evaluation criteria, the suggested sample size, T , for the SPEC model selection algorithm can be determined. The SPEC model selection algorithm has been applied for $T=5(5)80$. In the sequel, the value of T for which the SPEC selection method achieves the best performance according to the evaluation criteria used, is examined. Figure 3 shows a plot of the average T , suggested by the evaluation criteria, across the forecasting horizons. The bar charts are a graphical representation of the number of evaluation criteria by which the performance of the models selected by the SPEC algorithm were judged 'better' than the performance of any other single model (measured on the right hand side vertical axis). For a 16–36 days ahead forecasting horizon, the appropriate T , as concerns the specific data, ranges around 20 days with a standard deviation of 3.6 days. Table 4 provides more details for the sample size of the SPEC selection method suggested by the evaluation criteria and its standard deviation for both the entire 16–36 days ahead forecasting horizon and for each day individually. The SPEC model selection algorithm shows a better performance for a sample size of about 20 days.

Several results in the literature (e.g. Lopez and Walter, 2001; Christoffersen and Jacobs, 2003; Ferreira and Lopez, 2003) reveal that the simplest model specifications are chosen a disproportionately large percentage of the time, while others

(e.g. Vilasuso, 2002; Brooks and Persaud, 2003, Hansen and Lunde, 2003; Giot and Laurent, 2003, 2004; Angelidis *et al.*, 2004; Degiannakis, 2004) indicate that the more flexible an ARCH model is, the more adequate it is in volatility forecasting, compared to parsimonious models. In order to give the reader a sense of which of the 85 models was selected most often, Table 5 presents the models selected by the SPEC(20) algorithm. For example, the model with AR(0) conditional mean and GARCH(0,1) conditional variance was picked on 34 trading days. As concerns the conditional variance function, the GARCH, EGARCH and TARCH models were picked as the *most suitable* in the 38%, 39%, and 23% of the cases, respectively. On the basis of the results of Table 5, the SPEC algorithm does not appear to be noticeably biased towards selecting a specific type of model. This is in line with Degiannakis and Xekalaki's (2001) findings. Tables for the remaining sample sizes T of the SPEC algorithm were also constructed giving qualitatively similar results.¹²

VI. Comparison of the SPEC Criterion to Other Methods of Model Selection

Most of the methods used in the time series literature for selecting the appropriate model are based on evaluating the ability of the models to describe the data. Standard model selection criteria such as the Akaike information criterion [AIC] (Akaike, 1973) and the Schwarz Bayesian criterion [SBC] (Schwarz, 1978) have widely been used in the ARCH literature, despite the fact that their statistical

¹² These tables are available upon request.

Table 4. Average sample size for the SPEC model selection algorithm suggested by the evaluation criteria for both the entire 16–36 days ahead forecasting horizon and for each day individually

Forecasting horizon (in number of days ahead)	Average sample size suggested by the Evaluation Criteria rating the performance of the SPEC selection algorithm 'best'			Average sample size suggested by all the Evaluation Criteria considered		
	Number of criteria	Average sample size	Standard deviation	Number of criteria	Average sample size	Standard deviation
16–36	366	19.7	3.6	420	19.9	3.7
16	12	23.8	1.7	20	26.0	2.5
17	14	20.7	1.5	20	23.5	2.9
18	18	24.7	2.8	20	24.3	2.7
19	18	25.0	3.3	20	24.3	3.3
20	18	23.6	3.3	20	23.0	3.3
21	18	23.3	3.4	20	22.5	3.4
22	18	20.0	3.8	20	19.5	3.6
23	18	19.4	3.8	20	19.0	3.7
24	18	19.4	3.8	20	19.0	3.7
25	18	17.2	2.9	20	17.0	2.8
26	18	17.2	2.9	20	17.0	2.8
27	18	17.8	3.6	20	17.5	3.4
28	18	18.3	3.1	20	18.0	2.9
29	18	18.3	3.1	20	18.0	2.9
30	18	20.6	6.5	20	20.0	6.2
31	18	16.1	1.4	20	16.0	1.4
32	18	17.8	3.6	20	17.5	3.4
33	16	15.6	0.8	20	20.0	6.2
34	18	21.1	4.7	20	20.5	4.5
35	18	17.8	3.6	20	17.5	3.4
36	18	17.8	2.9	20	17.5	2.8

Table 5. Number of ARCH models selected by the SPEC(20) algorithm for 1081 trading days, classified by the types of models considered for their conditional means and variances

		Type of conditional mean model					Total
		AR(0)	AR(1)	AR(2)	AR(3)	AR(4)	
Type of conditional variance model	GARCH(0,1)	34	14	0	8	6	62
	GARCH(0,2)	12	0	2	6	1	21
	GARCH(1,1)	31	24	4	12	14	85
	GARCH(1,2)	38	9	0	9	16	72
	GARCH(2,1)	21	0	8	1	6	36
	GARCH(2,2)	62	21	4	7	41	135
	TARCH(0,1)	0	0	0	0	0	0
	TARCH(0,2)	0	0	2	4	2	8
	TARCH(1,1)	1	6	0	2	3	12
	TARCH(1,2)	2	61	20	3	19	105
	TARCH(2,1)	0	1	26	22	5	54
	TARCH(2,2)	23	13	19	5	10	70
	EGARCH(0,1)	81	31	4	51	14	181
	EGARCH(0,2)	6	8	6	0	5	25
	EGARCH(1,1)	4	20	13	8	11	56
	EGARCH(1,2)	14	28	21	10	14	87
EGARCH(2,1)	27	7	24	1	13	72	
Total		356	243	153	149	180	1081

properties in the ARCH context are unknown. Heq (1996), based on a set of Monte Carlo simulations, showed how the information criteria behave under the presence of ARCH effects. In small sample situations, the SBC is the best performing criterion. These are defined in terms of $l_n(\hat{\theta})$, the maximized value of the log-likelihood function of a model, $\hat{\theta}$ is the maximum likelihood estimator of the parameter vector θ based on a sample of size n and $\tilde{\theta}$ denotes the dimension of θ , thus:

$$AIC = l_n(\hat{\theta}) - \tilde{\theta} \quad (21)$$

$$SBC = l_n(\hat{\theta}) - 2^{-1}\tilde{\theta}\ln(n) \quad (22)$$

In addition, model selection is mainly based on the evaluation of some loss functions for each of the competing models. In this section, the statistical criteria, which were considered in Section IV as measures in evaluating the predictive performance of a variance model, are considered as criteria for the selection of ARCH models. In particular, the model selection methods presented in Table 6 are considered and their ability to predict future volatility is investigated.

Applying the SPEC algorithm, the sum of squared standardized one-step-ahead prediction errors, $\sum_{t=1}^T \hat{\varepsilon}_{t|t-1}^2 / \hat{\sigma}_{t|t-1}^2$, was estimated considering various values for T . Therefore, each of the model selection criteria, in Table 6, was computed considering various values for T , and, in particular, $T = 10(10)80$. The AIC and SBC criteria were computed based on the rolling sample of constant size equal to 500, or $n = 500$, that is used at each time to estimate the parameters of the models. Selecting a strategy for each method of model selection naturally amounts to selecting the model, which, at time k , has the lowest value of the formula is indicated in Table 6.

As concerns the AIC and SBC selection methods, they do not achieve the lowest value of the evaluation criteria in almost all the cases, which is indicative of the inability of the in-sample model selection methods to suggest the models with superior volatility forecasting performance. The general conclusion is that the loss functions presented in Table 6 do not lead to the selection of the ARCH processes which track closer the realized volatility. The HAAEVar, HASEVar and HASEDev methods show a better performance, as they select the ARCH models with the lowest value of the evaluation criteria, around the forecasting horizon ranging from 16 to 36 days ahead. So, they might be used in selecting that model that generates 'better' volatility predictions. The other selection methods failed to pick the models that perform 'better' in almost all the cases. Indicatively, Table 7 presents the percentage of

Table 6. Methods of selection of ARCH models. $\sigma_{t(N)}^2$ denotes the forecasting variance over an N day period measured at day t and $s_{t(N)}^2$ denotes the realized variance over the same period

1. Square Error of Conditional Variance (SEVar):

$$\sum_{t=1}^T \left((\sigma_{t(N)}^2 - s_{t(N)}^2)^2 \right) \quad (23)$$

2. Absolute Error of Conditional Variance (AEVar):

$$\sum_{t=1}^T \left(|\sigma_{t(N)}^2 - s_{t(N)}^2| \right) \quad (24)$$

3. Square Error of Conditional Standard Deviation (SEDev):

$$\sum_{t=1}^T \left((\sigma_{t(N)} - s_{t(N)})^2 \right) \quad (25)$$

4. Absolute Error of Conditional Standard Deviation (AEDDev):

$$\sum_{t=1}^T \left(|\sigma_{t(N)} - s_{t(N)}| \right) \quad (26)$$

5. Heteroscedasticity Adjusted Squared Error of Cond. Variance (HASEVar):

$$\sum_{t=1}^T \left(1 - \frac{s_{t(N)}^2}{\sigma_{t(N)}^2} \right)^2 \quad (27)$$

6. Heteroscedasticity Adjusted Absolute Error of Cond. Variance (HAAEVar):

$$\sum_{t=1}^T \left| 1 - \frac{s_{t(N)}^2}{\sigma_{t(N)}^2} \right| \quad (28)$$

7. Heteroscedasticity Adjusted Squared Error of Cond. St. Deviation (HASEDev):

$$\sum_{t=1}^T \left(1 - \frac{s_{t(N)}}{\sigma_{t(N)}} \right)^2 \quad (29)$$

8. Heteroscedasticity Adjusted Absolute Error of Cond. St. Deviation (HAAEDev):

$$\sum_{t=1}^T \left| 1 - \frac{s_{t(N)}}{\sigma_{t(N)}} \right| \quad (30)$$

9. Logarithmic Error of Conditional Variance (LEVar):

$$\sum_{t=1}^T \ln \left(\frac{s_{t(N)}^2}{\sigma_{t(N)}^2} \right)^2 \quad (31)$$

cases the models selected by the HAAEVar and LEVar model selection methods perform 'better' as judged by the evaluation criteria. The performance of the HASEVar and HASEDev selection methods is similar to that of the HAAEVar method, whereas the performance of the remaining methods is similar to that of the LEVar method. Full tables for all the methods considered are available upon request. In order to investigate whether the suggested model selection method indicates the ARCH models that track closer the realized volatility, the predictive ability of these loss functions must be compared to the volatility forecasting ability of the SPEC criterion, and mainly for a forecasting horizon ranging from 16 to 36 days ahead.

Of main interest is whether the ARCH models selected by the SPEC algorithm yield values for the evaluation criteria that are lower than those corresponding to the ARCH models selected by the model selection methods summarized in Table 6. As concerns forecasting horizons of 4–7 trading weeks ahead the performance of the SPEC algorithm is by far the best. Table 8 presents, indicatively, the percentage of times the ARCH models selected by the SPEC algorithm achieve lower values for the corresponding evaluation criteria and the specific forecasting horizons than the models selected by the HAAEVar and LEVar model selection methods. The SPEC algorithm performs 'better' than the other methods of model selection in about 90% of the cases. This percentage is lower when the SPEC algorithm is compared to the HAAEVar, HASEVar and HASEDev methods. Nevertheless, even in such cases, the opponent methods select the ARCH models that track closer future volatility much less frequently than the SPEC algorithm. The percentage of times, an opponent to the SPEC algorithm selects the most appropriate models in forecasting future volatility, is highest in the case of the HAAEVar method. However, only in the 23% of the cases, the ARCH models selected by the HAAEVar method perform 'better' than the models selected by the SPEC criterion, for any of the three horizon ranges. The performance of the remaining model selection methods is similar to that of the LEVar method. Full tables with the comparison of all the model selection methods to the SPEC algorithm are available upon request.

VII. Discussion

The SPEC method, for selecting an ARCH model among several competing models, amounts to

choosing the model with the lowest sum of squared standardized one-step-ahead forecasting errors. It incorporates the idea of 'jumping' from one model to another, as stock market behaviour alters. Thus, using the SPEC model selection algorithm every time a volatility forecast is required, allows shifting from the model used to predict the conditional variance the previous time to another.

In this paper, a number of evaluation criteria, for forecasting horizons ranging from one day to 100 days ahead, were applied and it was found that the ARCH models, picked by the SPEC model selection algorithm, generate 'better' predictions of the volatility. Thus, the SPEC selection method appears to be a useful tool in guiding choice of the appropriate model for estimating future volatility, with applications in evaluating portfolios, derivatives and financial risk.

Brooks and Persaud's (2003) evaluation approach was adopted and multistep-ahead forecasts were evaluated based on overlapping time periods. Alternatively, one might like to consider non-overlapping time periods and apply other evaluation schemes, such as those proposed by Diebold and Mariano (1995), Hansen and Lund (2003) or Hansen *et al.* (2003).

A topic worth exploring is the application of SPEC algorithm on models that account for recent developments in the area of volatility. Considering fractional integration of the conditional variance, for example, is an interesting question with regard to investigating SPEC's applicability further. (For more details, see, e.g., Giot and Laurent (2003) and Degiannakis (2004)). Finally, assessing the utility of the SPEC algorithm as a tool in model selection for ARCH models with non-normally distributed conditional innovations would be equally worthy as it would bring into play more general forms of models in the statistical and econometric literature.

References

- Adrangi, B. and Chatrath, A. (2003) Non-linear dynamics in futures prices: evidence from the coffee, sugar and cocoa exchange, *Applied Financial Economics*, **13**, 245–56.
- Akaike, H. (1973) Information theory and an extension of the maximum likelihood principle, *Proceedings of the Second International Symposium on Information Theory* (Eds) B. N. Petrov and F. Csaki, Akademiai Kiado, Budapest, pp. 267–81.
- Alizadeh, S., Brandt, M. W. and Diebold, F. X. (2002) Range-based estimation of stochastic volatility models, *Journal of Finance*, **LV11**, 1047–91.

- Andersen, T. and Bollerslev, T. (1997) Intraday periodicity and volatility persistence in financial markets, *Journal of Empirical Finance*, **4**, 115–58.
- Andersen, T. and Bollerslev, T. (1998a) Answering the skeptics: yes, standard volatility models do provide accurate forecasts, *International Economic Review*, **39**, 885–905.
- Andersen, T. and Bollerslev, T. (1998b) DM-dollar volatility: intraday activity patterns, macroeconomic announcements and longer-run dependencies, *Journal of Finance*, **53**, 219–65.
- Andersen, T., Bollerslev, T. and Lange, S. (1999) Forecasting financial market volatility: sample frequency vis-à-vis forecast horizon, *Journal of Empirical Finance*, **6**, 457–77.
- Andersen, T., Bollerslev, T. and Cai, J. (2000a) Intraday and interday volatility in the Japanese stock market, *Journal of International Financial Markets, Institutions and Money*, **10**, 107–30.
- Andersen, T., Bollerslev, T., Diebold, F. X. and Labys, P. (2000b) Exchange rate returns standardized by realized volatility are (nearly) Gaussian, *Multinational Finance Journal*, **4**, 159–79.
- Andersen, T., Bollerslev, T., Diebold, F. X. and Labys, P. (2001a) The distribution of exchange rate volatility, *Journal of the American Statistical Association*, **96**, 42–55.
- Andersen, T., Bollerslev, T., Diebold, F. X. and Ebens, H. (2001b) The distribution of stock return volatility, *Journal of Financial Economics*, **61**, 43–76.
- Andersen, T., Bollerslev, T., Diebold, F. X. and Labys, P. (2003) Modeling and forecasting realized volatility, *Econometrica*, **71**, 529–626.
- Andersen, T., Bollerslev, T. and Diebold, F. X. (2005) Parametric and nonparametric volatility measurement. *Handbook of Financial Econometrics* (Eds) Y. Aït-Sahalia and L. P. Hansen, Amsterdam, North Holland.
- Angelidis, T., Benos, A. and Degiannakis, S. (2004) The use of GARCH models in VaR estimation, *Statistical Methodology*, **1**, 105–28.
- Barkoulas, J. and Travlos, N. (1998) Chaos in an emerging capital market? The case of the Athens stock exchange, *Applied Financial Economics*, **8**, 231–43.
- Barkoulas, J., Baum, C. F. and Travlos, N. (2000) Long memory in the Greek stock market, *Applied Financial Economics*, **10**, 177–84.
- Barndorff-Nielsen, O. E. and Shephard, N. (1998) Aggregation and model construction for volatility models, Manuscript, University of Aarhus and Nuffield College, Oxford, Department of Mathematical Sciences.
- Basle Committee on Banking Supervision (1998). *International Convergence of Capital Measurement and Capital Standards*, Basle Committee on Banking Supervision, Switzerland.
- Bera, A. K. and Higgins, M. L. (1993) ARCH models: properties, estimation and testing, *Journal of Economic Surveys*, **7**, 305–66.
- Berndt, E. R., Hall, B. H., Hall, R. E. and Hausman, J. A. (1974) Estimation and inference in nonlinear structural models, *Annals of Economic and Social Measurement*, **3**, 653–65.
- Bollerslev, T. (1986) Generalized autoregressive conditional heteroskedasticity, *Journal of Econometrics*, **31**, 307–27.
- Bollerslev, T. and Wooldridge, J. M. (1992) Quasi-maximum likelihood estimation and inference in dynamic models with time-varying covariances, *Econometric Reviews*, **11**, 143–72.
- Bollerslev, T., Chou, R. and Kroner, K. F. (1992) ARCH modeling in finance: a review of the theory and empirical evidence, *Journal of Econometrics*, **52**, 5–59.
- Bollerslev, T., Engle, R. F. and Nelson, D. (1994) ARCH Models, in *Handbook of Econometrics*, Vol. 4 (Eds) R. Engle and D. McFadden, Elsevier Science, Amsterdam, pp. 2959–3038.
- Brock, W. (1986) Distinguishing random and deterministic systems: abridged version, *Journal of Economic Theory*, **40**, 168–95.
- Brock, W. A., Dechert, W. D. and Scheinkman, J. A. (1987). A test for independence based on the correlation dimension, Working Paper No. 8702, Department of Economics, University of Wisconsin, Madison, WI. SSRI.
- Brooks, C. and Persaud, G. (2003) The effect of asymmetries on stock index return value-at-risk estimates, *The Journal of Risk Finance*, **4**, 29–42.
- Campbell, J., Lo, A. and MacKinlay, A. C. (1997) *The Econometrics of Financial Markets*, Princeton University Press, New Jersey.
- Christoffersen, P. and Jacobs, K. (2003) Which volatility model for option evaluation? Manuscript, Faculty of Management, McGill University.
- Cohen, K., Hawawini, G., Maier, S., Schwartz, R. and Whitcomb, D. (1983) Friction in the trading process and the estimation of systematic risk, *Journal of Financial Economics*, **12**, 263–78.
- Degiannakis, S. (2004) Volatility forecasting: evidence from a fractional integrated asymmetric power arch skewed-*t* model, *Applied Financial Economics*, **14**, 1333–42.
- Degiannakis, S. and Xekalaki, E. (2001) Using a prediction error criterion for model selection in forecasting option prices, Technical Report No. 131, Department of Statistics, Athens University of Economics and Business. ([http://statathens.aueb.gr/~exek/papers/Xekalaki-TechnRep131\(2001\)ft.pdf](http://statathens.aueb.gr/~exek/papers/Xekalaki-TechnRep131(2001)ft.pdf))
- Degiannakis, S. and Xekalaki, E. (2004) Autoregressive conditional heteroscedasticity models: a review, *Quality Technology and Quantitative Management*, **1**, 271–324.
- Degiannakis, S. and Xekalaki, E. (2005) Predictability and model selection in the context of ARCH models, *Journal of Applied Stochastic Models in Business and Industry*, **21**, 55–82.
- Diebold, F. X. and Mariano, R. (1995) Comparing predictive accuracy, *Journal of Business and Economic Statistics*, **13**, 253–63.
- Dimson, E. (1979) Risk measurement when shares are subject to infrequent trading, *Journal of Financial Economics*, **7**, 197–226.
- Ebens, H. (1999) Realized stock volatility, Working Paper No. 420, Department of Economics, Johns Hopkins University.
- Engle, R. F., Hong, C. H., Kane, A. and Noh, J. (1993) Arbitrage valuation of variance forecasts with simulated options, *Advances in Futures and Options Research*, **6**, 393–415.

- Ferreira, M. A. and Lopez, J. A. (2003). Evaluating interest rate covariance models within a value-at-risk framework, Manuscript, Economic Research Department, Federal Reserve Bank of San Francisco.
- Franses, P. H. and Homelen, P. V. (1998) On forecasting exchange rates using neural networks, *Applied Financial Economics*, **8**, 589–96.
- Frey, R. and Michaud, P. (1997). The effect of GARCH-type volatilities on prices and payoff-distributions of derivative assets – a simulation study, Working Paper, ETH Zurich.
- Giot, P. and Laurent, S. (2003) Value-at-risk for long and short trading positions, *Journal of Applied Econometrics*, **18**, 641–64.
- Giot, P. and Laurent, S. (2004) Modelling daily value-at-risk using realized volatility and ARCH type models, *Journal of Empirical Finance*, **11**, 379–98.
- Glosten, L., Jagannathan, R. and Runkle, D. (1993) On the relation between the expected value and the volatility of the nominal excess return on stocks, *Journal of Finance*, **48**, 1779–801.
- González-Rivera, G., Lee, T.-H. and Mishra, S. (2004) Forecasting volatility: a reality check based on option pricing, utility function, value-at-risk and predictive likelihood, *International Journal of Forecasting*, **20**, 629–45.
- Gourieroux, C. (1997) *ARCH Models and Financial Applications*, Springer-Verlag, New York.
- Hamilton, J. (1994) *Time Series Analysis*, Princeton University Press, New Jersey.
- Hansen, P. R. and Lunde, A. (2003) A forecast comparison of volatility models: does anything beat a GARCH(1,1)? Working Paper, Brown University, Department of Economics.
- Hansen, P. R., Lunde, A. and Nason, J. M. (2003) Choosing the best volatility models: the model confidence set approach, Working Paper, Brown University, Department of Economics.
- Hecq, A. (1996) IGARCH effect on autoregressive lag length selection and causality tests, *Applied Economics Letters*, **3**, 317–23.
- Hertz, J., Krogh, A. and Palmer, R. (1991) *Introduction to the Theory of Neural Computation*, Addison-Wesley Publishing Company, Reading, MA.
- Heynen, R. and Kat, H. (1994) Volatility prediction: a comparison of the stochastic volatility, Garch(1,1) and Egarch(1,1) models, *Journal of Derivatives*, **94**, 50–65.
- Hol, E. and Koopman, S. (2000) Forecasting the variability of stock index returns with stochastic volatility models and implied volatility, Discussion Paper No. 104/4, Tinbergen Institute.
- Holden, A. (1986) *Chaos*, Princeton University Press, New Jersey.
- Hsieh, D. (1991) Chaos and nonlinear dynamics: application to financial markets, *Journal of Finance*, **46**, 1839–77.
- Hutchinson, J., Lo, A. and Poggio, T. (1994) A nonparametric approach to the pricing and hedging of derivative securities via learning networks, *Journal of Finance*, **49**, 851–89.
- Jasic, T. and Wood, D. (2004) The profitability of daily stock market indices trades based on neural network predictions: case study for the S&P500, the DAX, the TOPIX and the FTSE in the Period 1965–1999, *Applied Financial Economics*, **14**, 285–97.
- Kibble, W. F. (1941) A two variate gamma type distribution, *Sankhya*, **5**, 137–50.
- Klaassen, F. (2002) Improving GARCH volatility forecasts with regime-switching GARCH, in *Advances in Markov-Switching Models* (Eds) J. D. Hamilton and B. Raj, Physica Verlag, New York, pp. 223–54.
- Lo, A. and MacKinlay, A. C. (1988) Stock market prices do not follow random walks: evidence from a simple specification test, *Review of Financial Studies*, **1**, 41–66.
- Lo, A. and MacKinlay, A. C. (1990) An econometric analysis of non-synchronous trading, *Journal of Econometrics*, **45**, 181–212.
- Lopez, J. A. and Walter, C. A. (2001) Evaluating covariance matrix forecasts in a value-at-risk framework, *Journal of Risk*, **3**, 69–98.
- Marquardt, D. W. (1963) An algorithm for least squares estimation of nonlinear parameters, *Journal of the Society for Industrial and Applied Mathematics*, **11**, 431–41.
- Nelson, D. (1991) Conditional heteroskedasticity in asset returns: a new approach, *Econometrica*, **59**, 347–70.
- Pagan, A. R. and Schwert, G. W. (1990) Alternative models for conditional stock volatility, *Journal of Econometrics*, **45**, 267–90.
- Peel, D. A. and Speight, A. E. H. (1996) Is the US business cycle asymmetric? Some further evidence, *Applied Economics*, **28**, 405–15.
- Perez-Rodriguez, J. V., Torra, S. and Andrada-Felix, J. (2005) Are Spanish Ibex35 stock future index returns forecasted with non-linear models?, *Applied Financial Economics*, **15**, 963–75.
- Plasmans, J., Verkooijen, W. and Daniels, H. (1998) Estimating structural exchange rate models by artificial neural networks, *Applied Financial Economics*, **8**, 541–51.
- Poggio, T. and Girosi, F. (1990) Networks for approximation and learning, *Proceeding of the IEEE*, special issue: *Neural Networks I: Theory and Modeling*, **78**, 1481–97.
- Priestley, M. (1988) *Nonlinear and Non-Stationary Time Series Analysis*, Academic Press, San Diego.
- Sadorsky, P. (2005) Stochastic volatility forecasting and risk management, *Applied Financial Economics*, **15**, 121–35.
- Saez, M. (1997) Option pricing under stochastic volatility and stochastic interest rate in the Spanish case, *Applied Financial Economics*, **7**, 379–94.
- Saltoglu, B. (2003) Comparing forecasting ability of parametric and non-parametric methods: an application with Canadian monthly interest rates, *Applied Financial Economics*, **13**, 169–79.
- Scholes, M. and Williams, J. (1977) Estimating betas from non-synchronous data, *Journal of Financial Economics*, **5**, 309–28.
- Schwarz, G. (1978) Estimating the dimension of a model, *Annals of Statistics*, **6**, 461–4.
- Selcuk, F. (2005) Asymmetric stochastic volatility in emerging stock market, *Applied Financial Economics*, **15**, 867–74.
- Shephard, N. (1996) Statistical aspects of ARCH and stochastic volatility models, in *Time Series Models in Econometrics, Finance and Other Fields* (Eds) D. R. Cox, D. V. Hinkley, and O. E. Barndorff-Nielsen, Chapman & Hall, London, pp. 1–67.

- Taylor, S. J. (1994) Modelling stochastic volatility: a review and comparative study, *Mathematical Finance*, **4**, 183–204.
- Teräsvirta, T., Tjøstheim, D. and Granger, C. (1994) Aspects of modeling nonlinear time series, in *Handbook of Econometrics*, Vol. 4 (Eds) R. Engle and D. McFadden, Elsevier Science, Amsterdam, pp. 2919–57.
- Thompson, J. and Stewart, H. (1986) *Nonlinear Dynamics and Chaos*, John Wiley and Sons, NY.
- Tong, H. (1990) *Nonlinear Time Series: A Dynamic System Approach*, Oxford University Press, Oxford.
- Vilasuso, J. (2002) Forecasting exchange rate volatility, *Economics Letters*, **76**, 59–64.
- Walsh, D. M. and Tsou, G. Y.-G. (1998) Forecasting index volatility: sampling interval and non-trading effects, *Applied Financial Economics*, **8**, 477–85.
- West, K. D. and Cho, D. (1995) The predictive ability of several models of exchange rate volatility, *Journal of Econometrics*, **69**, 367–91.
- West, K. D., Edison, H. J. and Cho, D. (1993) A utility based comparison of some models for exchange rate volatility, *Journal of International Economics*, **35**, 23–45.
- White, H. (1992) *Artificial Neural Networks: Approximation and Learning Theory*, Blackwell Publishers, Cambridge, MA.
- Xekalaki, E. and Degiannakis, S. (2005). Evaluating volatility forecasts in option pricing in the context of a simulated options market, *Computational Statistics and Data Analysis*, Special Issue on Computational Econometrics, **49**, 611–29.
- Xekalaki E., Panaretos, J. and Psarakis, S. (2003) A predictive model evaluation and selection approach – the correlated gamma ratio distribution, *Stochastic Musings: Perspectives from the Pioneers of the Late 20th Century* (Ed.) J. Panaretos, Associates Publishers, Lawrence Erlbaum, pp. 188–202.
- Yu, J. (2002) Forecasting volatility in the New Zealand stock market, *Applied Financial Economics*, **12**, 193–202.