

examples

- 1990 USA general social survey, 284–285, 287, 290–291, 315–316
- AB/BA crossover trial, 322–325
- aircraft damage dataset, 245–249
- bioassay — factor 8 blood clotting times, 204–217
- bivariate simulated count data, 293, 296
- body temperature data, 21–24, 72–74
- English premiership football data for season 2006–2007, 249–255
- estimation of risk in medical studies using Monte Carlo, 33–34
- estimation of the prevalence of a disease, 18
- evaluation of candidate school tutors, 171–172
- goals of the Greek national team in Euro 2004, 15–18
- inverse Gaussian simulated dataset, 278
- Kobe Bryant's field goals in NBA, 19–20, 46–55, 125, 131–132, 134, 136–138, 141–143, 148, 313–315, 399–402
- Manchester United's goals, 354–356, 360, 366, 368–371, 376, 378
- odds ratios analysis from various studies, 319–320
- outstanding car insurance claim amounts, 347–353
- oxygen uptake experiment, 221–226
- relative risk using Bayes theorem, 4
- repeated measurements of blood pressure, 308–312
- schizotypal personality data, 178–184, 191–195, 329–331, 345–347
- senility symptoms data, 56–71, 77–81, 263–268
- simulated data
 - binomial, 272
 - bivariate Poisson, 293, 296
 - of Dellaportas et al. (2002), 414–419
 - inverse Gaussian, 278–279
 - normal data, 357–359, 361, 363–368, 371–374, 377
 - Poisson, 270
- slice sampling illustration, 77–81
- soft drink delivery times, 157–161, 164–167, 281–282, 383–386, 403–405, 424, 429–431
- treatment comparison in matched-pair clinical trials, 327–328
- water polo World Cup 2000 data, 334–336